

The NEW ENGLAND JOURNAL of MEDICINE

VOL. 352 NO. 12

ESTABLISHED IN 1812

MARCH 24, 2005

WWW.NEJM.ORG



1169 THIS WEEK IN THE JOURNAL

PERSPECTIVE

- 1171 Uncompensated Hospital Care J.S. Weissman
1173 Monitoring Risk in Approved Drugs S. Okie
1176 Herbal Medicine in Europe P.A.G.M. De Smet

ORIGINAL ARTICLES

- 1179 Addition of Clopidogrel to Aspirin and Fibrinolytic
Therapy for Myocardial Infarction
M.S. Sabatine and Others
1190 Hospitalization for Mental Illness among Parents
after the Death of a Child
J. Li and Others
1197 ACVBP versus CHOP plus Radiotherapy
for Localized Aggressive Lymphoma
F. Reyes and Others
1206 Day-Night Pattern of Sudden Death in Obstructive
Sleep Apnea
A.S. Gami and Others

CLINICAL PRACTICE

- 1215 Pertussis — Not Just for Kids
E.L. Hewlett and K.M. Edwards

REVIEW ARTICLE

- 1223 Medical Progress: Polycystic Ovary Syndrome
D.A. Ehrmann

IMAGES IN CLINICAL MEDICINE

- 1237 Renal Papillary Necrosis Unmasking
Sickle Cell Disease
M. Voulgarelis and P.D. Ziakas
e12 Severe Fecal Impaction
Y. Loubières and O. Chereau

CASE RECORDS OF THE MASSACHUSETTS GENERAL HOSPITAL

- 1238 A Man with Acute Respiratory Failure
D.M. Systrom and C. Wittram

EDITORIALS

- 1248 Concurrent Antiplatelet and Fibrinolytic Therapy
R.A. Lange and L.D. Hillis
1250 Stages of Aggressive Non-Hodgkin's Lymphoma
J.O. Armitage
1252 Financing Health Care
R. Kronick

SOUNDING BOARD

- 1255 Health Care Vouchers and Universal Coverage
E.J. Emanuel and V.R. Fuchs
1260 Do We Really Want Broad Access to Health Care?
J.J. Mongan and T.H. Lee

1264 CORRESPONDENCE

- Palifermin and Chemotherapy-Induced Oral Mucositis
Care of the Potential Organ Donor
Contagious Acute Gastrointestinal Infections
Cancer of the Ovary
Postmenopausal Bleeding and an Ovarian Mass
Why "Why" Matters
Nosocomial Transmission of Cryptococcosis

1273 BOOK REVIEWS

1276 CORRECTIONS

1277 CONTINUING MEDICAL EDUCATION

Owned & published by the MASSACHUSETTS MEDICAL SOCIETY © 2005.
All rights reserved. ISSN 0028-4793.

YOUR CUSTOMER NUMBER: 000810276

FOLEY HOAG LIBRARY

2109
167
988
7

MAR 23 2005



ORIGINAL ARTICLE

Day–Night Pattern of Sudden Death in Obstructive Sleep Apnea

Apoor S. Gami, M.D., Daniel E. Howard, B.S., Eric J. Olson, M.D.,
and Virend K. Somers, M.D., Ph.D.

ABSTRACT

BACKGROUND

The risk of sudden death from cardiac causes in the general population peaks from 6 a.m. to noon and has a nadir from midnight to 6 a.m. Obstructive sleep apnea is highly prevalent and associated with neurohormonal and electrophysiological abnormalities that may increase the risk of sudden death from cardiac causes, especially during sleep.

METHODS

We reviewed polysomnograms and the death certificates of 112 Minnesota residents who had undergone polysomnography and had died suddenly from cardiac causes between July 1987 and July 2003. For four intervals of the day, we compared the rates of sudden death from cardiac causes among people with obstructive sleep apnea and the following: the rates among people without obstructive sleep apnea, the rates in the general population, and the expectations according to chance. For each interval, we assessed the median apnea–hypopnea index and the relative risk of sudden death from cardiac causes. We similarly analyzed sudden death from cardiac causes during three time intervals that correlate with usual sleep–wake cycles.

RESULTS

From midnight to 6 a.m., sudden death from cardiac causes occurred in 46 percent of people with obstructive sleep apnea, as compared with 21 percent of people without obstructive sleep apnea ($P=0.01$), 16 percent of the general population ($P<0.001$), and the 25 percent expected by chance ($P<0.001$). People with sudden death from cardiac causes from midnight to 6 a.m. had a significantly higher apnea–hypopnea index than those with sudden death from cardiac causes during other intervals, and the apnea–hypopnea index correlated directly with the relative risk of sudden death from cardiac causes from midnight to 6 a.m. For people with obstructive sleep apnea, the relative risk of sudden death from cardiac causes from midnight to 6 a.m. was 2.57 (95 percent confidence interval, 1.87 to 3.52). The analysis of usual sleep–wake cycles showed similar results.

CONCLUSIONS

People with obstructive sleep apnea have a peak in sudden death from cardiac causes during the sleeping hours, which contrasts strikingly with the nadir of sudden death from cardiac causes during this period in people without obstructive sleep apnea and in the general population.

From the Divisions of Cardiovascular Diseases (A.S.G., V.K.S.), Pulmonary and Critical Care Medicine (E.J.O.), and Hypertension (V.K.S.), Department of Internal Medicine (A.S.G., D.E.H., E.J.O., V.K.S.), Mayo Clinic College of Medicine, Rochester, Minn. Address reprint requests to Dr. Somers at the Division of Cardiovascular Diseases, Mayo Clinic College of Medicine, 200 First St. SW, Rochester, MN 55905, or at somers.virend@mayo.edu.

N Engl J Med 2005;352:1206-14.

Copyright © 2005 Massachusetts Medical Society.

THE RISK OF SUDDEN DEATH FROM CARDIAC causes in the general population is significantly greater during the morning hours after waking (i.e., from 6 a.m. to noon) than during the other six-hour intervals of the day.¹ Also, there is a marked nadir in the risk of sudden death from cardiac causes during sleep (i.e., from midnight to 6 a.m.).¹ This same day-night pattern has been shown for the incidence of sudden death from cardiac causes in patients with heart failure² and for the incidence of acute myocardial infarction.¹ The increase in risk in the morning may in part be due to changes in sympathetic activity,³ baroreflex sensitivity,⁴ coagulability,⁵ and electrophysiological abnormalities⁶⁻⁸ during the waking morning hours, all of which may predispose persons to cardiac ischemia and fatal arrhythmias.

Obstructive sleep apnea affects 17 to 24 percent of North American adults.⁹ In people with obstructive sleep apnea, nocturnal repetitive episodes of acute apnea elicit hypoxemia, hypercapnia, increased sympathetic drive,¹⁰ surges in blood pressure,¹⁰ increases in cardiac-wall stress,¹¹ and cardiac arrhythmias.¹²⁻¹⁴ Obstructive sleep apnea is also associated with hypercoagulability, vascular oxidative stress, systemic inflammation, and endothelial dysfunction.¹⁵ Thus, people with obstructive sleep apnea have severe perturbations of autonomic, hemodynamic, humoral, and vascular regulation during sleep that contrast with the physiology of normal sleep. It is not known whether these nocturnal abnormalities are associated with an increased risk of sudden death from cardiac causes during the night, a time when the risk of sudden death from cardiac causes is at its lowest level for the general population.¹

We tested the hypothesis that people with obstructive sleep apnea are more likely to have sudden death from cardiac causes from midnight to 6 a.m. than during the other six-hour intervals of the day. We also tested the hypothesis that the frequency of sudden death from cardiac causes from midnight to 6 a.m. is higher in people with obstructive sleep apnea than in people without obstructive sleep apnea, the general population, and what is expected by chance (i.e., 25 percent per six-hour interval).

METHODS

STUDY POPULATION

We included subjects who authorized their records to be used for research, and the institutional review

board of the Mayo Foundation approved the study. We identified Minnesota residents who underwent diagnostic polysomnography at the Mayo Clinic Sleep Disorders Center from July 1, 1987, to July 31, 2003, and confirmed their vital status (available for 99.6 percent) with the use of records from the Minnesota Department of Health and the Mayo Clinic. We then manually reviewed the deceased persons' death certificates (available for 83.8 percent) and included in our study those who died of sudden death from cardiac causes before July 31, 2003 (112 people).

CLASSIFICATION OF SUDDEN DEATH FROM CARDIAC CAUSES

Manual review of death certificates and data from the Minnesota Department of Health provided the immediate and underlying causes of death, time of death, and time interval from onset of symptoms to death. A state nosologist used the methods of the National Center for Health Statistics to assign causes of death. We classified the following causes of death as sudden death from cardiac causes: "sudden cardiac death," "cardiac dysrhythmia," "cardiac arrhythmia," "cardiac arrest," "cardiorespiratory arrest," or "coronary heart disease" or "myocardial infarction" in which the interval from symptoms to death was an hour or less.

Persons were excluded if the time of sudden death from cardiac causes was not stated or if the death certificate provided information that explicitly contradicted the following definition of sudden death from cardiac causes: "natural death due to cardiac causes, heralded by abrupt loss of consciousness within one hour of the onset of acute symptoms; preexisting heart disease may have been known to be present but the time and mode of death are unexpected"¹⁶ and nontraumatic.¹⁶ The occurrence of sudden death from cardiac causes during sleep was an exception to the criteria for "loss of consciousness within one hour of the acute onset of symptoms,"¹⁶ since some instances of sudden death from cardiac causes occurring during sleep may preclude awakening and the development of symptoms, or the witnessing of these events by others. The death certificates of six people stated that sudden death from cardiac causes was unwitnessed or presumed, but their time of death was noted, and these people were included in the study. The death certificates of all other persons did not state whether sudden death from cardiac causes was witnessed or unwitnessed. Thirty-six death

certificates (32 percent) provided specific information about the time interval from symptoms to death.

CLASSIFICATION OF OBSTRUCTIVE SLEEP APNEA

The sleep evaluations of all 112 persons were conducted by a sleep specialist at the Mayo Clinic Sleep Disorder Center. We reviewed each person's first diagnostic polysomnogram, which included measures of the electroencephalogram, electro-oculogram, electromyogram, electrocardiogram, thoracoabdominal excursions, pulse oximetry, and naso-oral airflow.

An apnea was defined as cessation of airflow for at least 10 seconds in the presence of thoracoabdominal ventilatory efforts, and a hypopnea as a reduction in airflow of at least 30 percent with a decrease in oxygen saturation of 2 percent or more for at least 10 seconds in the presence of thoracoabdominal ventilatory efforts. The apnea-hypopnea index was calculated as the sum of apneas and hypopneas per hour of sleep. According to American Academy of Sleep Medicine criteria, an apnea-hypopnea index of 5 or more established the diagnosis of obstructive sleep apnea.¹⁷

We collected data on the demographics, coexisting conditions, height, and weight of each person at the time of polysomnography. We considered that continuous positive airway pressure had been used if it was prescribed after the sleep study (which was noted by the sleep-medicine physician reviewing the study results and plan of care) and if a subsequent note in the medical record stated that continuous positive airway pressure was being used. The personnel who collected data from polysomnograms and confirmed sleep diagnoses were different from those who collected data from death certificates and confirmed death diagnoses, and the procedures were performed in a masked fashion.

STATISTICAL ANALYSIS

Characteristics of the study population were described as means \pm SD and counts (with percentages) and were compared according to the status of obstructive sleep apnea and the time of sudden death from cardiac causes with the use of the two-tailed t-test, Fisher's exact test, and analysis of variance. The frequency distribution of sudden death from cardiac causes for the four six-hour intervals was compared between persons with and persons without obstructive sleep apnea with the use of

Fisher's exact test. The proportion of people with obstructive sleep apnea who had sudden death from cardiac causes during each time interval was compared, with the use of the binomial distribution, with the proportion of people who had sudden death from cardiac causes during the same time interval in a historical control population (from a meta-analysis of 19,390 persons with sudden death from cardiac causes in the general population)¹ and with the proportion that was expected by chance to have sudden death from cardiac causes during each quarter of the day (i.e., 25 percent).

The apnea-hypopnea indexes of persons who had sudden death from cardiac causes during each time interval were expressed as medians (with interquartile ranges) and compared with the use of the Kruskal-Wallis test. For persons with and without obstructive sleep apnea, we calculated the relative risk and 95 percent confidence interval of sudden death from cardiac causes during each 6-hour interval as compared with the remaining 18 hours of the day.

We also performed similar analyses of sudden death from cardiac causes in persons with and persons without obstructive sleep apnea for three eight-hour intervals of the day (6 a.m. to 2 p.m., 2 p.m. to 10 p.m., and 10 p.m. to 6 a.m.). These intervals better represent usual sleep-wake cycles.

RESULTS

PATIENT CHARACTERISTICS

Characteristics of the study population are described in Table 1. The diagnoses for the 34 persons without obstructive sleep apnea were no sleep disorder (53 percent), central sleep apnea (23 percent), periodic leg movement disorder (9 percent), obesity-hypoventilation syndrome (6 percent), narcolepsy (6 percent), and hypoventilation due to neuromuscular disease (3 percent). There was a higher proportion of men in the group with obstructive sleep apnea than in the group without obstructive sleep apnea, but the two groups were similar in terms of age, body-mass index, and coexisting conditions. There were no differences in characteristics, or reported use of continuous positive airway pressure, between persons with sudden death from cardiac causes during the various intervals of the day. The use of continuous positive airway pressure was reported in 15 percent of patients with an apnea-hypopnea index of 5 to 19, 39 percent of patients with an apnea-hypopnea index of 20 to 39, and 58

Table 1. Characteristics of the Study Population at the Time of Polysomnography, According to the Status of Obstructive Sleep Apnea (OSA) and the Time of Sudden Death from Cardiac Causes.*

Characteristic	OSA Status			Time of Death				P Value†
	OSA (N=78)	No OSA (N=34)	P Value	Midnight–5:59 a.m. (N=43)	6 a.m.–11:59 a.m. (N=30)	Noon–5:59 p.m. (N=16)	6 p.m.–11:59 p.m. (N=23)	
Age — yr	70±10	67±13	0.18	67±11	71±13	71±11	71±8	0.47
Male sex — no. (%)	64 (82)	21 (62)	0.02	33 (77)	24 (80)	11 (69)	17 (74)	0.85
Body-mass index‡	34±7	33±9	0.71	35±8	32±7	33±11	33±5	0.59
Hypertension — no. (%)	51 (65)	22 (65)	0.94	27 (63)	20 (67)	8 (50)	18 (78)	0.32
Diabetes mellitus — no. (%)	32 (41)	12 (35)	0.57	20 (47)	6 (20)	9 (56)	9 (39)	0.06
Coronary artery disease — no. (%)	54 (69)	19 (56)	0.29	27 (63)	18 (60)	10 (63)	18 (78)	0.52
Congestive heart failure — no. (%)	43 (55)	18 (53)	0.83	24 (56)	13 (43)	10 (63)	14 (61)	0.51
Cerebrovascular disease — no. (%)	17 (22)	7 (21)	0.89	7 (16)	8 (27)	3 (19)	6 (26)	0.68
Dysrhythmia — no. (%)§	14 (18)	3 (9)	0.22	6 (14)	3 (10)	5 (31)	3 (13)	0.27

* Plus-minus values are means ±SD.

† The P value for time of death is for the comparison of groups according to time of death.

‡ The body-mass index is the weight in kilograms divided by the square of the height in meters.

§ A diagnosis of dysrhythmia includes a history of ventricular tachycardia, ventricular fibrillation, an implanted cardiac defibrillator, or resuscitation from cardiac arrest.


percent of patients with an apnea-hypopnea index of 40 or more ($P < 0.001$).

TIME DISTRIBUTION OF SUDDEN DEATH FROM CARDIAC CAUSES


Figure 1 shows the primary results. From midnight to 6 a.m., the frequency of sudden death from cardiac causes was significantly higher in persons with obstructive sleep apnea than in persons without obstructive sleep apnea (46 percent vs. 21 percent, $P = 0.01$), than in the general population (46 percent vs. 16 percent, $P < 0.001$), and than was expected by chance (46 percent vs. 25 percent, $P < 0.001$). From 6 a.m. to noon, the frequency of sudden death from cardiac causes was significantly lower in persons with obstructive sleep apnea than in persons without obstructive sleep apnea (20 percent vs. 41 percent, $P = 0.04$). From noon to 6 p.m., the frequency of sudden death from cardiac causes was significantly lower in persons with obstructive sleep apnea than in persons without obstructive sleep apnea (9 percent vs. 26 percent, $P = 0.02$), than in the general population (9 percent vs. 29 percent, $P < 0.001$), and than was expected by chance (9 percent vs. 25 percent, $P < 0.001$).

SEVERITY OF OBSTRUCTIVE SLEEP APNEA

Persons with sudden death from cardiac causes from midnight to 6 a.m. had a significantly higher



median apnea-hypopnea index than those with sudden death from cardiac causes from 6 a.m. to noon (39 vs. 8, $P < 0.001$) and noon to 6 p.m. (39 vs. 11, $P < 0.001$), and a similar median apnea-hypopnea



index to that in persons with sudden death from cardiac causes from 6 p.m. to midnight (39 vs. 41) (Fig. 2A). As compared with persons without obstructive sleep apnea, the relative risk of sudden death from cardiac causes from midnight to 6 a.m. was

1.87 (95 percent confidence interval, 0.86 to 4.04) for persons with mild-to-moderate obstructive sleep apnea (apnea-hypopnea index, 5 to 39) and 2.61 (95 percent confidence interval, 1.27 to 5.38) for persons with severe obstructive sleep apnea (apnea-hypopnea index, ≥ 40) (Fig. 2B).

RELATIVE RISK OF SUDDEN DEATH FROM CARDIAC CAUSES

For persons with obstructive sleep apnea, the relative risk of sudden death from cardiac causes from midnight to 6 a.m. (as compared with the remaining 18 hours of the day) was 2.57 (95 percent confidence interval, 1.87 to 3.52), from 6 a.m. to noon was 0.77 (95 percent confidence interval, 0.49 to 1.21), from noon to 6 p.m. was 0.30 (95 percent confidence interval, 0.15 to 0.61), and from 6 p.m. to midnight was 0.97 (95 percent confidence interval, 0.64 to 1.46). For persons without obstructive sleep apnea, the relative risk of sudden death from cardiac causes from midnight to 6 a.m. (as compared with the remaining 18 hours of the day) was 0.77 (95 percent confidence interval, 0.36 to 1.66), from 6 a.m. to noon was 2.10 (95 percent confidence interval, 1.14 to 3.85), from noon to 6 p.m. was 1.08 (95 percent confidence interval, 0.54 to 2.16), and from 6 p.m. to midnight was 0.40 (95 percent confidence interval, 0.15 to 1.08) (Fig. 3).

SLEEP-WAKE CYCLES

The sleep-wake cycles are described in Figure 4. Included are the day-night pattern of sudden death from cardiac causes (Fig. 4A), the severity of obstructive sleep apnea (Fig. 4B), and the relative risk of sudden death from cardiac causes during the three time intervals that represent usual sleep-wake cycles (Fig. 4C).

DISCUSSION

This study shows that persons with obstructive sleep apnea have a significantly increased risk of sudden death from cardiac causes during the sleeping hours, which is in striking contrast to the nadir of sudden death from cardiac causes during this time in persons without obstructive sleep apnea and in the general population. In the analysis of sudden death from cardiac causes from midnight to 6 a.m. (which allows for the comparison of our results to results of large studies in the general population) and in the analysis of sudden death from cardiac causes from 10 p.m. to 6 a.m. (which is more rele-

vant to sleep-related pathophysiology), a marked nocturnal peak in sudden death from cardiac causes was observed in persons with obstructive sleep apnea. In more than half of persons with obstructive sleep apnea, sudden death from cardiac causes occurred between 10 p.m. and 6 a.m. By contrast, the persons without obstructive sleep apnea in our study sample had a day-night pattern of sudden death from cardiac causes very similar to that in the general population, with a peak in sudden death from cardiac causes from 6 a.m. to noon.

Furthermore, our analysis showed that the severity of obstructive sleep apnea correlated directly with the risk of nocturnal sudden death from cardiac causes, such that the relative risk of sudden death from cardiac causes during the sleeping hours was 40 percent higher in persons with severe obstructive sleep apnea (apnea-hypopnea index, ≥ 40) than in those with mild-to-moderate obstructive sleep apnea (apnea-hypopnea index, 5 to 39).

The mean age at which sudden death from cardiac causes occurred in persons with obstructive sleep apnea was similar to the mean age of sudden death from cardiac causes in persons without obstructive sleep apnea and in the general population.¹⁸ Our findings should not be generalized to younger persons with obstructive sleep apnea. Also, this finding suggests that obstructive sleep apnea does not hasten sudden death from cardiac causes; however, our study cannot address whether obstructive sleep apnea increases the overall risk of sudden death from cardiac causes.

Our data may also provide insights into the conventional understanding of the day-night pattern of sudden death from cardiac causes in the general population. First, obstructive sleep apnea may be implicated in some of the 16 percent of cases of sudden death from cardiac causes that occur between midnight and 6 a.m. in the general population,¹ since obstructive sleep apnea affects at least 25 million adult Americans⁹ and remains undiagnosed in the vast majority of these people.¹⁹ Second, the risk of sudden death from cardiac causes from 6 a.m. to noon in persons without obstructive sleep apnea may be even greater than the risk that is currently supported by data from the general population, since those data include people with and people without obstructive sleep apnea, and we showed that the presence of obstructive sleep apnea is associated with a lower risk during this period.

The marked nocturnal nadir of sudden death from cardiac causes in the general population prob-




Figure 3. Relative Risk of Sudden Death from Cardiac Causes during Six-Hour Intervals as Compared with the Remaining 18 Hours of the Day, for 78 Persons with and 34 Persons without Obstructive Sleep Apnea (OSA).

The squares represent the point estimates of relative risk, and the I bars the 95 percent confidence intervals.

ably reflects the normal physiology of sleep, in which sympathetic activity is decreased³ and cardiac dysrhythmias are uncommon.²⁰ The unique day-night pattern of sudden death from cardiac causes that we observed in persons with obstructive sleep apnea is consistent with their exposure during sleep to critical mechanisms that could promote sudden death from cardiac causes.

Obstructive sleep apnea is characterized by repetitive collapse of the pharyngeal airway during sleep, resulting in complete or partial cessation of airflow, sometimes hundreds of times nightly. The resultant hypoxemia can lead to nocturnal cardiac ischemia²¹ and ventricular arrhythmias.²² Apneic episodes elicit increased sympathetic activity,¹⁰ blood-pressure elevations,¹⁰ and platelet aggregation.²³ Obstructive sleep apnea is associated with abnormalities in cardiac autonomic and electrophysiological factors, including heart rate variability,²⁴ the duration of the QT interval,²⁵ baroreflex function, and chemoreceptor sensitivity.²⁶ Serious and potentially fatal arrhythmias occur during sleep in patients with obstructive sleep apnea^{12-14,22,27-31} and are attenuated by effective treatment.^{14,29-32} Thus, multiple pathophysiological mechanisms occur during sleep in persons with obstructive sleep apnea and may explain an increased risk of nocturnal sudden death from cardiac causes.


Two previous studies evaluated the day-night

Figure 4. Sudden Death from Cardiac Causes According to Usual Sleep–Wake Cycles.

Panel A shows day–night patterns of sudden death from cardiac causes on the basis of eight-hour time intervals for 78 persons with and 34 persons without obstructive sleep apnea. Panel B shows the apnea–hypopnea index for persons with sudden death from cardiac causes during eight-hour intervals. The line within each box represents the median apnea–hypopnea index, and the box represents the interquartile range (25th percentile to 75th percentile). The figure includes data from persons with and from persons without obstructive sleep apnea ($P=0.001$) for the comparison of the apnea–hypopnea index according to the time of sudden death. Panel C shows the relative risk of sudden death from cardiac causes during 8-hour intervals, as compared with the remaining 16 hours of the day, for 78 persons with and 34 persons without obstructive sleep apnea. The squares represent the relative risk point estimates, and the I bars the 95 percent confidence intervals.

pattern of sudden death from cardiac causes in persons with sleep disorders.^{33,34} Neither study specifically examined obstructive sleep apnea. The first study obtained histories of snoring from the cohabitants of 321 men who had sudden death from cardiac causes. It found that habitual snorers were more likely than occasional snorers or nonsnorers to have sudden death from cardiac causes during sleep and between 4 a.m. and 8 a.m.³³ The second study reported that among 13 persons with sleep-disordered breathing (not just obstructive sleep apnea) and sudden death from cardiac causes, none died during sleep.³⁴ In contrast, our study assessed the day–night pattern of sudden death from cardiac causes in a large group of nonselected persons with or without confirmed obstructive sleep apnea as determined by the gold standard of diagnostic tests.

One limitation of the present study relates to the recognized difficulty of establishing the diagnosis and timing of sudden death from cardiac causes. Our use of data from death certificates to identify sudden death from cardiac causes has precedent in previous epidemiologic studies^{35,36} and has been validated in large populations,^{37–40} including our regional population.^{39,40} Furthermore, we used information from the death certificates about the time interval from the onset of symptoms to death in order to corroborate the diagnosis of sudden death from cardiac causes on the basis of current definitions.¹⁶ For example, we were able to exclude persons who were diagnosed with sudden death from cardiac causes on the basis of old criteria that allowed a 24-hour interval between symptoms and death.



Our methods also improve on past studies that searched databases of diagnostic codes to identify potential cases, since we identified persons with sudden death from cardiac causes by our direct review of every available death record from the larger population. Qualifiers such as “witnessed,” “unwitnessed,” and “presumed” were infrequently noted on the death certificates. Therefore, we included all cases of sudden death from cardiac causes regardless of these statements. Thus, persons with sudden

death from cardiac causes during sleep, which may preclude the development of symptoms or the witnessing of the event by others, were exceptions to the criteria that required timing of symptoms and a change in consciousness, and they were included in the study when a time of death was noted. It is important to note that we collected sleep data and death data independently of one another, and in a masked fashion, in order to avoid diagnostic suspicion bias.

Another limitation of the study relates to the ascertainment and interpretation of treatment with continuous positive airway pressure. During the years of this study, continuous-positive-airway-pressure devices that stored the time of use were rarely used. The medical records lacked objective data regarding frequency, duration, or effectiveness of continuous positive airway pressure. Therefore, we could not verify whether continuous positive airway pressure was used in the days before sudden death from cardiac causes or during the event.

That continuous positive airway pressure may have been generally ineffective is suggested by the finding that the risk of nocturnal sudden death

from cardiac causes increased with increasing severity of obstructive sleep apnea despite more reported use of continuous positive airway pressure with increasing severity of obstructive sleep apnea. Because of the lack of objective data from continuous-positive-airway-pressure devices, subjective data regarding their use were necessary. This is an important limitation of retrospective assessments of treatment of obstructive sleep apnea.

This study shows that persons with obstructive sleep apnea have a significant alteration in the well-established day-night pattern of sudden death from cardiac causes evident in the general population. Our study cannot address the question of whether obstructive sleep apnea increases the overall risk of sudden death from cardiac causes. Given our findings and the relatively high prevalence of obstructive sleep apnea in Western populations, this is an important question that remains to be answered.

Supported by grants (HL61560, HL65176, HL73211, and M01 R-00585) from the National Institutes of Health, by the Dana Foundation, and by the Mayo Clinic College of Medicine. Dr. Gami is also supported by the Dr. Ralph and Marian C. Falk Medical Research Trust Fellowship for Clinical Research.

REFERENCES

- Cohen MC, Rohitla KM, Lavery CE, Muller JE, Mittleman MA. Meta-analysis of the morning excess of acute myocardial infarction and sudden cardiac death. *Am J Cardiol* 1997;79:1512-6. [Erratum, *Am J Cardiol* 1998;81:260.]
- Moser DK, Stevenson WG, Woo MA, Stevenson LW. Timing of sudden death in patients with heart failure. *J Am Coll Cardiol* 1994;24:963-7.
- Somers V, Dyken ME, Mark AL, Abboud FM. Sympathetic-nerve activity during sleep in normal subjects. *N Engl J Med* 1993;328:303-7.
- Legramante JM, Marciani MG, Placidi F, et al. Sleep-related changes in baroreflex sensitivity and cardiovascular autonomic modulation. *J Hypertens* 2003;21:1555-61.
- Andreotti F, Davies GJ, Hackett DR, et al. Major circadian fluctuations in fibrinolytic factors and possible relevance to time of onset of myocardial infarction, sudden cardiac death and stroke. *Am J Cardiol* 1988;62:635-7.
- Bexton RS, Vallin HO, Camm AJ. Diurnal variation of the QT interval — influence of the autonomic nervous system. *Br Heart J* 1986;55:253-8.
- Huikuri HV, Niemela MJ, Ojala S, Rantala A, Ikaheimo MJ, Airaksinen KE. Circadian rhythms of frequency domain measures of heart rate variability in healthy subjects and patients with coronary artery disease: effects of arousal and upright posture. *Circulation* 1994;90:121-6.
- Huikuri HH, Yli-Mayry S, Linnaluoto MK, Ikaheimo MJ. Diurnal fluctuations in human ventricular and atrial refractoriness. *Pacing Clin Electrophysiol* 1995;18:1362-8.
- Young T, Peppard PE, Gottlieb DJ. Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 2002;165:1217-39.
- Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Invest* 1995;96:1897-904.
- Virolainen J, Ventila M, Turto H, Kupari M. Effect of negative intrathoracic pressure on left ventricular pressure dynamics and relaxation. *J Appl Physiol* 1995;79:455-60.
- Guilleminault C, Connolly SJ, Winkle RA. Cardiac arrhythmia and conduction disturbances during sleep in 400 patients with sleep apnea syndrome. *Am J Cardiol* 1983;52:490-4.
- Fichter J, Bauer D, Arampatzis S, Fries R, Heisel A, Sybrecht GW. Sleep-related breathing disorders are associated with ventricular arrhythmias in patients with an implantable cardioverter-defibrillator. *Chest* 2002;122:558-61.
- Harbison J, O'Reilly P, McNicholas WT. Cardiac rhythm disturbances in the obstructive sleep apnea syndrome: effects of nasal continuous positive airway pressure therapy. *Chest* 2000;118:591-5.
- Shamsuzzaman AS, Gersh BJ, Somers VK. Obstructive sleep apnea: implications for cardiac and vascular disease. *JAMA* 2003;290:1906-14.
- Priori SG, Aliot E, Blomstrom-Lundqvist C, et al. Task Force on Sudden Cardiac Death of the European Society of Cardiology. *Eur Heart J* 2001;22:1374-450. [Erratum, *Eur Heart J* 2002;23:257.]
- Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research: the report of the American Academy of Sleep Medicine Task Force. *Sleep* 1999;22:667-89.
- Fox CS, Evans JC, Larson MG, Kannel WB, Levy D. Temporal trends in coronary heart disease mortality and sudden cardiac death from 1950 to 1999: the Framingham Heart Study. *Circulation* 2004;110:522-7.
- Young T, Evans L, Finn L, Palta M. Estimation of the clinically diagnosed proportion of sleep apnea syndrome in middle-aged men and women. *Sleep* 1997;20:705-6.
- Clarke JM, Hamer J, Shelton JR, Taylor S, Venning GR. The rhythm of the normal human heart. *Lancet* 1976;1:508-12.
- Schafer H, Koehler U, Ploch T, Peter JH.

- Sleep-related myocardial ischemia and sleep structure in patients with obstructive sleep apnea and coronary heart disease. *Chest* 1997;111:387-93.
22. Shepard JW Jr, Garrison MW, Grither DA, Dolan GF. Relationship of ventricular ectopy to oxyhemoglobin desaturation in patients with obstructive sleep apnea. *Chest* 1985;88:335-40.
 23. Sanner BM, Konermann M, Tepel M, Groetz J, Mummenhoff C, Zidek W. Platelet function in patients with obstructive sleep apnoea syndrome. *Eur Respir J* 2000;16:648-52.
 24. Narkiewicz K, Montano N, Cogliati C, van de Borne PJ, Dyken ME, Somers VK. Altered cardiovascular variability in obstructive sleep apnea. *Circulation* 1998;98:1071-7.
 25. Gillis AM, Stoohs R, Guilleminault C. Changes in the QT interval during obstructive sleep apnea. *Sleep* 1991;14:346-50.
 26. Narkiewicz K, van de Borne PJ, Pesek CA, Dyken ME, Montano N, Somers VK. Selective potentiation of peripheral chemoreflex sensitivity in obstructive sleep apnea. *Circulation* 1999;99:1183-9.
 27. Zwillich C, Devlin T, White D, Douglas N, Weil J, Martin R. Bradycardia during sleep apnea: characteristics and mechanism. *J Clin Invest* 1982;69:1286-92.
 28. Fries R, Bauer D, Heisel A, et al. Clinical significance of sleep-related breathing disorders in patients with implantable cardioverter defibrillators. *Pacing Clin Electrophysiol* 1999;22:223-7.
 29. Koehler U, Fus E, Grimm W, et al. Heart block in patients with obstructive sleep apnoea: pathogenetic factors and effects of treatment. *Eur Respir J* 1998;11:434-9.
 30. Grimm W, Hoffmann J, Menz V, et al. Electrophysiologic evaluation of sinus node function and atrioventricular conduction in patients with prolonged ventricular asystole during obstructive sleep apnea. *Am J Cardiol* 1996;77:1310-4.
 31. Becker H, Brandenburg U, Peter JH, Von Wichert P. Reversal of sinus arrest and atrioventricular conduction block in patients with sleep apnea during nasal continuous positive airway pressure. *Am J Respir Crit Care Med* 1995;151:215-8.
 32. Tilkian AG, Guilleminault C, Schroeder JS, Lehrman KL, Simmons FB, Dement WC. Sleep-induced apnea syndrome: prevalence of cardiac arrhythmias and their reversal after tracheostomy. *Am J Med* 1977;63:348-58.
 33. Seppala T, Partinen M, Penttila A, Aspholm R, Tiainen E, Kaukianen A. Sudden death and sleeping history among Finnish men. *J Intern Med* 1991;229:23-8.
 34. Gonzalez-Rothi RJ, Foresman GE, Block AJ. Do patients with sleep apnea die in their sleep? *Chest* 1988;94:531-8.
 35. Muller JE, Ludmer PL, Willich SN, et al. Circadian variation in the frequency of sudden cardiac death. *Circulation* 1987;75:131-8.
 36. Goraya TY, Jacobsen SJ, Kottke TE, Frye RL, Weston SA, Roger VL. Coronary heart disease death and sudden cardiac death: a 20-year population-based study. *Am J Epidemiol* 2003;157:763-70.
 37. Iribarren C, Crow RS, Hannan PJ, Jacobs DR Jr, Luepker RV. Validation of death certificate diagnosis of out-of-hospital sudden cardiac death. *Am J Cardiol* 1998;82:50-3.
 38. Coady SA, Sorlie PD, Cooper LS, Folsom AR, Rosamond WD, Conwill DE. Validation of death certificate diagnosis for coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) Study. *J Clin Epidemiol* 2001;54:40-50.
 39. Folsom AR, Gomez-Marin O, Gillum RF, Kottke TE, Lohman W, Jacobs DR Jr. Out-of-hospital coronary death in an urban population — validation of death certificate diagnosis: the Minnesota Heart Survey. *Am J Epidemiol* 1987;125:1012-8.
 40. Goraya TY, Jacobsen SJ, Belau PG, Weston SA, Kottke TE, Roger VL. Validation of death certificate diagnosis of out-of-hospital coronary heart disease deaths in Olmsted County, Minnesota. *Mayo Clin Proc* 2000;75:681-7.

Copyright © 2005 Massachusetts Medical Society.

CLINICAL TRIAL REGISTRATION

The *Journal* encourages investigators to register their clinical trials in a public trials registry. The members of the International Committee of Medical Journal Editors plan to consider clinical trials for publication only if they have been registered (see *N Engl J Med* 2004;351:1250-1). The National Library of Medicine's www.clinicaltrials.gov is a free registry, open to all investigators, that meets the committee's requirements.